

Please check the examination details below before entering your candidate information

Candidate surname		Other names	
Centre Number		Candidate Number	

Pearson Edexcel Level 3 GCE

Tuesday 6 June 2023

Afternoon (Time: 2 hours)

Paper reference **9MA0/01**

Mathematics

Advanced

PAPER 1: Pure Mathematics 1

You must have:
Mathematical Formulae and Statistical Tables (Green), calculator

Total Marks

Candidates may use any calculator allowed by Pearson regulations. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B).
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
– *there may be more space than you need.*
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Inexact answers should be given to three significant figures unless otherwise stated.

Information

- A booklet 'Mathematical Formulae and Statistical Tables' is provided.
- There are 15 questions in this question paper. The total mark for this paper is 100.
- The marks for **each** question are shown in brackets
– *use this as a guide as to how much time to spend on each question.*

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ►

P72804A

©2023 Pearson Education Ltd.
N:1/1/1/1/1/




Pearson

1. Find

$$\int \frac{x^{\frac{1}{2}}(2x-5)}{3} dx$$

writing each term in simplest form.

(4)

$$\int \frac{1}{3} x^{\frac{1}{2}} (2x-5) dx$$

$$\int \frac{2}{3} x^{\frac{3}{2}} - \frac{5}{3} x^{\frac{1}{2}} dx$$

$$\frac{\frac{2}{3} x^{\frac{5}{2}}}{\frac{5}{2}} - \frac{\frac{5}{3} x^{\frac{3}{2}}}{\frac{3}{2}} + C$$

$$\frac{4}{15} x^{\frac{5}{2}} - \frac{10}{9} x^{\frac{3}{2}} + C$$



DO NOT WRITE IN THIS AREA

Question 1 continued

Lined area for writing answers.

(Total for Question 1 is 4 marks)



2.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

$$f(x) = 4x^3 + 5x^2 - 10x + 4a \quad x \in \mathbb{R}$$

where a is a positive constant.Given $(x - a)$ is a factor of $f(x)$,

$$f(a) = 0$$

(a) show that

$$a(4a^2 + 5a - 6) = 0 \quad (2)$$

(b) Hence

(i) find the value of a

(ii) use algebra to find the exact solutions of the equation

$$f(x) = 3 \quad (4)$$

$$a/ \quad 4a^3 + 5a^2 - 10a + 4a = 0$$

$$4a^3 + 5a^2 - 6a = 0$$

$$a(4a^2 + 5a - 6) = 0$$

$$b.i/ \quad a = 0 \quad a = \frac{3}{4} \quad a = -2$$

$$a = \frac{3}{4} \quad \text{as } a \text{ is a +ve constant.}$$

$$ii/ \quad 4x^3 + 5x^2 - 10x + 4\left(\frac{3}{4}\right) = 3$$

$$4x^3 + 5x^2 - 10x + 3 = 3$$

$$4x^3 + 5x^2 - 10x = 0$$

$$x(4x^2 + 5x - 10) = 0$$

$$x = 0, \quad \frac{-5 + \sqrt{185}}{8}, \quad \frac{-5 - \sqrt{185}}{8}$$



DO NOT WRITE IN THIS AREA

Question 2 continued

Handwriting practice area with horizontal lines.

(Total for Question 2 is 6 marks)



3. Relative to a fixed origin O

- the point A has position vector $5\mathbf{i} + 3\mathbf{j} + 2\mathbf{k}$
- the point B has position vector $2\mathbf{i} + 4\mathbf{j} + a\mathbf{k}$

where a is a positive integer.

(a) Show that $|\vec{OA}| = \sqrt{38}$

(1)

(b) Find the smallest value of a for which

$$|\vec{OB}| > |\vec{OA}|$$

(2)

$$a/ \quad \sqrt{5^2 + 3^2 + 2^2} = \underline{\underline{\sqrt{38}}}$$

$$b/ \quad \sqrt{2^2 + 4^2 + a^2} > \sqrt{38}$$

$$2^2 + 4^2 + a^2 > 38$$

$$20 + a^2 > 38$$

$$a^2 > 18$$

$$\underline{\underline{a = 5}}$$



DO NOT WRITE IN THIS AREA

Question 3 continued

Lined area for writing the answer to Question 3.

(Total for Question 3 is 3 marks)



4.

In this question you must show all stages of your working.

Solutions relying entirely on calculator technology are not acceptable.

The curve C has equation $y = f(x)$ where $x \in \mathbb{R}$

Given that

- $f'(x) = 2x + \frac{1}{2} \cos x$
- the curve has a stationary point with x coordinate α
- α is small

$$f'(\alpha) = 0$$

- (a) use the small angle approximation for $\cos x$ to estimate the value of α to 3 decimal places.

$$\cos \theta \approx \underline{\underline{1 - \frac{\theta^2}{2}}} \quad (3)$$

The point $P(0, 3)$ lies on C

- (b) Find the equation of the tangent to the curve at P , giving your answer in the form $y = mx + c$, where m and c are constants to be found.

(2)

$$a/ \quad 2\alpha + \frac{1}{2} \cos \alpha = 0$$

$$2\alpha + \frac{1}{2} \left(1 - \frac{\alpha^2}{2}\right) = 0$$

$$2\alpha + \frac{1}{2} - \frac{\alpha^2}{4} = 0$$

$$-\frac{1}{4} \alpha^2 + 2\alpha + \frac{1}{2} = 0$$

$$\alpha = \cancel{8.24}$$

$$\alpha = \underline{\underline{-0.243}}$$

$$b/ \quad f'(x) = 2x + \frac{1}{2} \cos x \quad (0, 3)$$

$$m = 2(0) + \frac{1}{2} \cos(0)$$

$$= \frac{1}{2}$$

$$\underline{\underline{y = \frac{1}{2}x + 3}}$$



DO NOT WRITE IN THIS AREA

Question 4 continued

Handwriting practice area with horizontal lines.

(Total for Question 4 is 5 marks)



5. A continuous curve has equation $y = f(x)$.

The table shows corresponding values of x and y for this curve, where a and b are constants.

x	3	3.2	3.4	3.6	3.8	4
y	a	16.8	b	20.2	18.7	13.5

The trapezium rule is used, with all the y values in the table, to find an approximate area under the curve between $x = 3$ and $x = 4$

Given that this area is 17.59

(a) show that $a + 2b = 51$

(3)

Given also that the sum of all the y values in the table is 97.2

(b) find the value of a and the value of b

(3)

$$a) \quad 0.2 \left(\frac{a}{2} + 16.8 + b + 20.2 + 18.7 + \frac{13.5}{2} \right) = 17.59$$

$$\frac{a}{2} + b + 62.45 = 87.95$$

$$\frac{a}{2} + b = \frac{51}{2}$$

$$a + 2b = 51$$

$$b) \quad a + b + 69.2 = 97.2$$

$$a + b = 28$$

$$a + 2b = 51$$

$$\underline{\underline{a = 5}}$$

$$\underline{\underline{b = 23}}$$



DO NOT WRITE IN THIS AREA

Question 5 continued

Handwriting practice area with horizontal lines.

(Total for Question 5 is 6 marks)



6.

$a = \log_2 x$

$b = \log_2(x + 8)$

Express in terms of a and/or b

$$(a) \log_2 \sqrt{x} \quad (1)$$

$$(b) \log_2(x^2 + 8x) \quad (2)$$

$$(c) \log_2\left(8 + \frac{64}{x}\right) \quad (3)$$

Give your answer in simplest form.

$$a/ \log_2 x^{\frac{1}{2}}$$

$$\frac{1}{2} \log_2 x = \underline{\underline{\frac{1}{2} a}}$$

$$b/ \log_2(x^2 + 8x)$$

$$\log_2(x(x+8))$$

$$\log_2 x + \log_2(x+8)$$

$$\underline{\underline{a + b}}$$

$$c/ \log_2\left(\frac{8}{1} + \frac{64}{x}\right)$$

$$\log_2\left(\frac{8x}{x} + \frac{64}{x}\right)$$

$$\log_2 \frac{8x + 64}{x}$$

$$\log_2(8x + 64) - \log_2 x$$

$$\log_2(8(x+8)) - \log_2 x$$

$$\log_2 8 + \log_2(x+8) - \log_2 x$$



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 6 continued

$$\underline{\underline{3 + b - a}}$$

(Total for Question 6 is 6 marks)



7. The function f is defined by

$$f(x) = 3 + \sqrt{x-2} \quad x \in \mathbb{R} \quad x > 2$$

(a) State the range of f (1)

(b) Find f^{-1} (3)

The function g is defined by

$$g(x) = \frac{15}{x-3} \quad x \in \mathbb{R} \quad x \neq 3$$

(c) Find $gf(6)$ (2)

(d) Find the exact value of the constant a for which

$$f(a^2 + 2) = g(a) \quad (2)$$

$$a/ \quad f(x) > 3$$

$$b/ \quad y = 3 + \sqrt{x-2}$$

$$y - 3 = \sqrt{x-2}$$

$$(y-3)^2 = x-2$$

$$x = (y-3)^2 + 2$$

$$f^{-1}(x) = (x-3)^2 + 2, \quad x > 3$$

$$c/ \quad f(6) = 3 + \sqrt{6-2}$$

$$= 5$$

$$g(5) = \frac{15}{5-3} = \frac{15}{2}$$



Question 7 continued

$$3 + \sqrt{a^2 + 2} - 2 = \frac{15}{a-3}$$

$$3 + \sqrt{a^2} = \frac{15}{a-3}$$

$$3 + a = \frac{15}{a-3}$$

$$(a+3)(a-3) = 15$$

$$a^2 - 9 = 15$$

$$a^2 = 24$$

$$a = \pm\sqrt{24}$$

reject $-\sqrt{24}$

$$\therefore a = \sqrt{24}$$

$$= \underline{\underline{2\sqrt{6}}}$$



Question 7 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 7 continued

Lined area for writing the answer to Question 7.

(Total for Question 7 is 8 marks)



8.


Diagram NOT
accurately drawn

Figure 1 shows the plan view of a stage.

The plan view shows two congruent triangles ABO and GFO joined to a sector $OCDEO$ of a circle, centre O , where

- angle $COE = 2.3$ radians
- arc length $CDE = 27.6$ m
- AOG is a straight line of length 15 m

(a) Show that $OC = 12$ m.

(2)

(b) Show that the size of angle AOB is 0.421 radians correct to 3 decimal places.

(2)

Given that the total length of the front of the stage, $BCDEF$, is 35 m,

(c) find the total area of the stage, giving your answer to the nearest square metre.

(6)

$$a/ \text{ Arc length} = r\theta$$

$$27.6 = r(2.3)$$

$$r = \frac{27.6}{2.3}$$

$$= 12 \text{ m}$$

$$b/ \frac{\pi - 2.3}{2} = \underline{\underline{0.421}}$$



Question 8 continued



$$c/ \quad 2x + 27.6 = 35$$

$$x = 3.7$$

$$OB = OC + x$$

$$= 12 + 3.7$$

$$= 15.7$$



$$\begin{aligned} \text{Triangles area} &= 2 \times \frac{1}{2} (7.5)(15.7) \sin(0.421) \\ &= 48.12 \end{aligned}$$

$$\text{Sector area} = \frac{1}{2} r^2 \theta$$

$$= \frac{1}{2} (12)^2 (2.3)$$

$$= 165.6$$

$$\text{Total} = 165.6 + 48.12 = \underline{\underline{214 \text{ m}^2}}$$



Question 8 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 8 continued

Handwriting practice area with horizontal lines.

(Total for Question 8 is 10 marks)



9. The first three terms of a geometric sequence are

$$3k + 4 \quad 12 - 3k \quad k + 16$$

where k is a constant.

(a) Show that k satisfies the equation

$$3k^2 - 62k + 40 = 0 \quad (2)$$

Given that the sequence converges, $-1 < r < 1$

(b) (i) find the value of k , giving a reason for your answer,

(ii) find the value of S_{∞} (5)

$$a/ \quad \frac{12 - 3k}{3k + 4} = \frac{k + 16}{12 - 3k}$$

$$(12 - 3k)(12 - 3k) = (k + 16)(3k + 4)$$

$$144 - 72k + 9k^2 = 3k^2 + 4k + 48k + 64$$

$$6k^2 - 124k + 80 = 0$$

$$\underline{3k^2 - 62k + 40 = 0}$$

$$b/ \quad k = 20 \quad \text{or} \quad \frac{2}{3}$$

$$\text{if } k = 20 \quad 64, -48, 36 \quad r = -\frac{3}{4}$$

$$\text{if } k = \frac{2}{3} \quad 6, 10, 16\frac{2}{3} \quad r = \frac{5}{3}$$

$$\underline{k = 20} \quad \text{because } -1 < r < 1 \quad \therefore r = -\frac{3}{4}$$



Question 9 continued

$$\begin{aligned} c/ \quad S_{\infty} &= \frac{a}{1-r} \\ &= \frac{64}{1 + \frac{3}{4}} \\ &= \frac{256}{7} \\ &= \underline{\underline{\quad\quad\quad}} \end{aligned}$$

(Total for Question 9 is 7 marks)



$$(x-a)^2 + (y-b)^2 = r^2$$

10. A circle C has equation


$$x^2 + y^2 + 6kx - 2ky + 7 = 0$$

where k is a constant.

(a) Find in terms of k ,

(i) the coordinates of the centre of C

(ii) the radius of C



(3)

The line with equation $y = 2x - 1$ intersects C at 2 distinct points.

(b) Find the range of possible values of k .

$$b^2 - 4ac > 0$$

(6)

$$a/ \quad x^2 + 6kx + y^2 - 2ky + 7 = 0$$

$$(x + 3k)^2 - 9k^2 + (y - k)^2 - k^2 + 7 = 0$$

$$(x + 3k)^2 + (y - k)^2 = 10k^2 - 7$$

$$i/ \quad \text{centre: } (-3k, k)$$

$$ii/ \quad \text{radius: } \sqrt{10k^2 - 7}$$

$$b/ \quad x^2 + 6kx + (2x-1)^2 - 2k(2x-1) + 7 = 0$$

$$\underline{x^2} + \underline{6kx} + \underline{4x^2} - \underline{4x} + \underline{1} - \underline{4kx} + \underline{2k} + \underline{7} = 0$$

$$5x^2 + 2kx - 4x + 2k + 8 = 0$$

$$5x^2 + (2k - 4)x + (2k + 8) = 0$$

$$b^2 - 4ac > 0$$

$$(2k - 4)^2 - 4(5)(2k + 8) > 0$$



Question 10 continued

$$4k^2 - 16k + 16 - 40k - 160 > 0$$

$$4k^2 - 56k - 144 > 0$$

$$\underline{k < 7 - \sqrt{85}} \quad \text{or} \quad \underline{k > 7 + \sqrt{85}}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



Question 10 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 10 continued

Handwriting practice area with horizontal lines.

(Total for Question 10 is 9 marks)



11.




Figure 2

The value, V pounds, of a mobile phone, t months after it was bought, is modelled by

$$V = ab^t$$

where a and b are constants.

Figure 2 shows the linear relationship between $\log_{10} V$ and t .

The line passes through the points $(0, 3)$ and $(10, 2.79)$ as shown.

Using these points,

(a) find the initial value of the phone, (2)

(b) find a complete equation for V in terms of t , giving the exact value of a and giving the value of b to 3 significant figures. (3)

Exactly 24 years after it was bought, the value of the phone was £320

(c) Use this information to evaluate the reliability of the model. (2)

a/ $\log_{10} V = 3$

$$V = 10^3$$

$$= 1000$$

$$\underline{\underline{\pounds 1000}}$$

b/ $m = \frac{2.79 - 3}{10} = -0.021$



Question 11 continued

$$\log_{10} v = -0.021t + 3$$

$$v = 10^{-0.021t + 3}$$

$$= 10^3 \times 10^{-0.021t}$$

$$= 1000 (10^{-0.021})^t$$

$$= 1000 (0.953)^t$$

$$c/ \quad v = 1000 (0.953)^{24}$$

$$= \underline{\underline{315}}$$

315 is close to 320 $\therefore$ the model seems to be reliable.



Question 11 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 11 continued

Lined area for writing the answer to Question 11.

(Total for Question 11 is 7 marks)



12.

$$y = \sin x$$

where x is measured in radians.

Use differentiation from first principles to show that

$$\frac{dy}{dx} = \cos x$$

You may

- use without proof the formula for $\sin(A \pm B)$
- assume that as $h \rightarrow 0$, $\frac{\sin h}{h} \rightarrow 1$ and $\frac{\cos h - 1}{h} \rightarrow 0$

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \quad (5)$$

$$= \lim_{h \rightarrow 0} \frac{\sin(x+h) - \sin x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin x \cos h + \cos x \sin h - \sin x}{h}$$

$$= \lim_{h \rightarrow 0} \frac{\sin x \cos h - \sin x}{h} + \frac{\cos x \sin h}{h}$$

$$\lim_{h \rightarrow 0} \sin x \left(\frac{\cos h - 1}{h} \right) + \cos x \left(\frac{\sin h}{h} \right)$$

$$\text{as } h \rightarrow 0 \quad f'(x) = \sin x (0) + \cos x (1)$$

$$= \underline{\underline{\cos x}}$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 12 continued

Handwriting practice area with horizontal lines.

(Total for Question 12 is 5 marks)



13. On a roller coaster ride, passengers travel in carriages around a track.


On the ride, carriages complete multiple circuits of the track such that

- the maximum vertical height of a carriage above the ground is 60 m
- a carriage starts a circuit at a vertical height of 2 m above the ground
- the ground is horizontal

The vertical height, H m, of a carriage above the ground, t seconds after the carriage starts the first circuit, is modelled by the equation

$$H = a - b(t - 20)^2$$

where a and b are positive constants.



(a) Find a complete equation for the model.

(b) Use the model to determine the height of the carriage above the ground when $t = 40$

(1)

In an alternative model, the vertical height, H m, of a carriage above the ground, t seconds after the carriage starts the first circuit, is given by

$$H = 29 \cos(9t + \alpha)^\circ + \beta \quad 0 \leq \alpha < 360^\circ$$

where α and β are constants.

(c) Find a complete equation for the alternative model.

(2)

Given that the carriage moves continuously for 2 minutes,

(d) give a reason why the alternative model would be more appropriate.

(1)

a/ Max height when $t = 20$

$$\therefore a = 60$$

$$\text{when } t = 0 \quad H = 2$$

$$2 = 60 - b(-20)^2$$

$$-58 = -b(400)$$

$$b = \frac{58}{400} = 0.145$$



Question 13 continued

$$H = 60 - 0.145(t - 20)^2$$

b/ when $t = 40$

$$H = 60 - 0.145(40 - 20)^2$$

$$= \underline{2 \text{ m}}$$

c/ $H = 29 \cos(9t + \alpha) + 31$


$$2 = 29 \cos \alpha + 31$$

$$\frac{-29}{29} = \cos \alpha$$

$$\alpha = 180^\circ$$

$$H = 29 \cos(9t + 180) + 31$$

d/ The alternative model will repeat



but the original will keep going down.



Question 13 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



DO NOT WRITE IN THIS AREA

Question 13 continued

Handwriting practice area with horizontal lines.

(Total for Question 13 is 7 marks)



14. Prove, using algebra, that

$$(n+1)^3 - n^3$$

is odd for all $n \in \mathbb{N}$

(4)

if n is even let $n = 2m$

$$(2m+1)^3 - (2m)^3$$

$$(2m+1)(2m+1)(2m+1) - 8m^3$$

$$(4m^2 + 4m + 1)(2m+1) - 8m^3$$

$$\underline{8m^3} + \underline{4m^2 + 8m^2 + 4m + 2m + 1} - \underline{8m^3}$$

$$12m^2 + 6m + 1$$

$$2(6m^2 + 3m) + 1 \quad \text{odd,}$$

even + 1 is odd

if n is odd let $n = 2m + 1$

$$(2m+1+1)^3 - (2m+1)^3$$

$$(2m+2)^3 - (8m^3 + 12m^2 + 6m + 1)$$

$$(2m+2)(2m+2)(2m+2) - 8m^3 - 12m^2 - 6m - 1$$

$$(4m^2 + 8m + 4)(2m+2) - 8m^3 - 12m^2 - 6m - 1$$

$$\underline{8m^3} + \underline{8m^2 + 16m^2 + 16m + 8m + 8} - \underline{8m^3} - \underline{12m^2} - \underline{6m} - \underline{1}$$

$$12m^2 + 18m + 7$$



Question 14 continued

$$12n^2 + 18n + 6 + 1$$

$$2(6n^2 + 9n + 3) + 1 \quad \text{odd.}$$

even + 1 is odd

$$\therefore \underline{(n+1)^3 - n^3} \quad \text{is odd}$$

for all $n \in \mathbb{N}$

(Total for Question 14 is 4 marks)

15. A curve has equation $y = f(x)$, where

$$f(x) = \frac{7xe^x}{\sqrt{e^{3x} - 2}} \quad x > \ln \sqrt[3]{2}$$

(a) Show that

$$f'(x) = \frac{7e^x(e^{3x}(2-x) + Ax + B)}{2(e^{3x} - 2)^{\frac{3}{2}}}$$

where A and B are constants to be found.

(5)

(b) Hence show that the x coordinates of the turning points of the curve are solutions of the equation

$$x = \frac{2e^{3x} - 4}{e^{3x} + 4}$$

(2)

The equation $x = \frac{2e^{3x} - 4}{e^{3x} + 4}$ has two positive roots α and β where $\beta > \alpha$

A student uses the iteration formula

$$x_{n+1} = \frac{2e^{3x_n} - 4}{e^{3x_n} + 4}$$

in an attempt to find approximations for α and β

Diagram 1 shows a plot of part of the curve with equation $y = \frac{2e^{3x} - 4}{e^{3x} + 4}$ and part of the line with equation $y = x$

Using Diagram 1 on page 42

(c) draw a staircase diagram to show that the iteration formula starting with $x_1 = 1$ can be used to find an approximation for β

(1)

Use the iteration formula with $x_1 = 1$, to find, to 3 decimal places,

(d) (i) the value of x_2

(ii) the value of β

(3)

Using a suitable interval and a suitable function that should be stated

(e) show that $\alpha = 0.432$ to 3 decimal places.

(2)



$$\frac{dy}{dx} = v \frac{du}{dx} + u \frac{dv}{dx}$$

Question 15 continued

$$u = 7xe^x$$

$$v = (e^{3x} - 2)^{\frac{1}{2}}$$

$$\frac{7x e^x}{7 e^x}$$

$$\frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$$

$$\frac{du}{dx} = 7e^x + 7xe^x$$

$$\frac{dv}{dx} = \frac{1}{2}(e^{3x} - 2)^{-\frac{1}{2}} \cdot 3e^{3x}$$

$$= \frac{3}{2}e^{3x}(e^{3x} - 2)^{-\frac{1}{2}}$$

$$f'(x) = \frac{(e^{3x} - 2)^{\frac{1}{2}}(7e^x + 7xe^x) - 7xe^x\left(\frac{3}{2}e^{3x}(e^{3x} - 2)^{-\frac{1}{2}}\right)}{e^{3x} - 2}$$

$$= \frac{(e^{3x} - 2)(7e^x + 7xe^x) - 7xe^x \cdot \frac{3}{2}e^{3x}}{(e^{3x} - 2)^{3/2}}$$

$$= \frac{2(e^{3x} - 2)(7e^x + 7xe^x) - 21xe^x \cdot e^{3x}}{2(e^{3x} - 2)^{3/2}}$$

$$= \frac{14e^x(1+x)(e^{3x} - 2) - 21xe^x \cdot e^{3x}}{2(e^{3x} - 2)^{3/2}}$$

$$= \frac{7e^x(2(1+x)(e^{3x} - 2) - 3xe^{3x})}{2(e^{3x} - 2)^{3/2}}$$

$$= \frac{7e^x(2(e^{3x} - 2 + xe^{3x} - 2x) - 3xe^{3x})}{2(e^{3x} - 2)^{3/2}}$$

Question 15 continued

Only use the copy of Diagram 1 if you need to redraw your answer to part (c).





Diagram 1



copy of Diagram 1

$$= \frac{7e^x (2(e^{3x} - 2 + xe^{3x} - 2x) - 3xe^{3x})}{2(e^{3x} - 2)^{3/2}}$$

$$= \frac{7e^x (2e^{3x} - 4 + 2xe^{3x} - 4x - 3xe^{3x})}{2(e^{3x} - 2)^{3/2}}$$

$$= \frac{7e^x (2e^{3x} - xe^{3x} - 4x - 4)}{2(e^{3x} - 2)^{3/2}}$$

$$= \frac{7e^x (e^{3x}(2-x) - 4x - 4)}{2(e^{3x} - 2)^{3/2}}$$

$$b) \frac{7e^x (e^{3x}(2-x) - 4x - 4)}{2(e^{3x} - 2)^{3/2}} = 0$$

$$\underline{7e^x} (e^{3x}(2-x) - 4x - 4) = 0$$

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Question 15 continued

$$e^{3x}(2-x) - 4x - 4 = 0$$

$$2e^{3x} - xe^{3x} - 4x - 4 = 0$$

$$2e^{3x} - 4 = xe^{3x} + 4x$$

$$2e^{3x} - 4 = x(e^{3x} + 4)$$

$$x = \frac{2e^{3x} - 4}{e^{3x} + 4}$$

d i/ $x_1 = 1$

$$x_2 = 1.502$$

ii/ $\beta = 1.968$

e/ $f'(x) = \frac{7e^x(e^{3x}(2-x) - 4x - 4)}{2(e^{3x} - 2)^{3/2}}$

$0.4315 \quad 0.4325$

$$f'(0.4315) = -5.789 \times 10^{-3}$$

$$f'(0.4325) = 0.0183$$

change of sign and continuous function

$\therefore \alpha = 0.432$ to 3dp.



Question 15 continued

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(Total for Question 15 is 13 marks)

TOTAL FOR PAPER IS 100 MARKS

